

Trying 3106016892...Open

Welcome to STN International! Enter x:x
LOGINID:sssptal653jxl
PASSWORD:
TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Dec 17 The CA Lexicon available in the CAPLUS and CA files
NEWS 3 Feb 06 Engineering Information Encompass files have new names
NEWS 4 Feb 16 TOXLINE no longer being updated
NEWS 5 Apr 23 Search Derwent WPINDEX by chemical structure
NEWS 6 Apr 23 PRE-1967 REFERENCES NOW SEARCHABLE IN CAPLUS AND CA
NEWS 7 May 07 DGENE Reload

NEWS EXPRESS April 18 CURRENT WINDOWS VERSION IS V6.0,
CURRENT MACINTOSH VERSION IS V5.0C (ENG) AND V5.0JB (JP),
AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2001
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 20:47:44 ON 22 MAY 2001

=> file biosis caplus medline scisearch

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.15	0.15

FILE 'BIOSIS' ENTERED AT 20:48:11 ON 22 MAY 2001
COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'CAPLUS' ENTERED AT 20:48:11 ON 22 MAY 2001
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'MEDLINE' ENTERED AT 20:48:11 ON 22 MAY 2001

FILE 'SCISEARCH' ENTERED AT 20:48:11 ON 22 MAY 2001
COPYRIGHT (C) 2001 Institute for Scientific Information (ISI) (R)

=> s cell denisty and image analysis

L1 0 CELL DENISTY AND IMAGE ANALYSIS

=> s cell denisty and classification

L2 0 CELL DENISTY AND CLASSIFICATION

=> .
=> s cell and classification and image analysis and tissue

3 FILES SEARCHED...
L3 196 CELL AND CLASSIFICATION AND IMAGE ANALYSIS AND TISSUE

=> s l3 and density

L4 22 L3 AND DENSITY

=> dup rem l4

PROCESSING COMPLETED FOR L4
L5 16 DUP REM L4 (6 DUPLICATES REMOVED)

=> d l5 bib ab 1-16

L5 ANSWER 1 OF 16 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 1
AN 2001:216929 BIOSIS
DN PREV200100216929
TI Microvascular development and growth of uterine **tissue** during
the estrous cycle in mares.
AU Ferreira-Dias, Graca M. (1); Serrao, Paula M. (1); Durao, Jose F. Costa
(1); Silva, Jose Robalo (1)
CS (1) Centro de Investigacao Interdisciplinar em Sanidade Animal, Faculdade
de Medicina Veterinaria, Rua Prof. Cid dos Santos Polo Universitario Alto
da Ajuda, 1300-477, Lisbon Portugal
SO American Journal of Veterinary Research, (April, 2001) Vol. 62, No. 4, pp.
526-530. print.
ISSN: 0002-9645.
DT Article
LA English
SL English
AB Objective-To document uterine growth and microvascular development in the
endometrium of uteri with differing degrees of fibrosis as well as uterine
growth throughout the estrous cycle of mares. Animals-30 mares.
Procedure-Uterine **tissue** was obtained during the breeding season
from a slaughter facility. Stage of estrous cycle of the mares was
assessed on the basis of ovarian structures and plasma progesterone
concentrations. Endometrium was characterized by use of light microscopy,
and blood vessel walls were marked by histochemical techniques.
Microvascular development was evaluated by a computerized **image**
analysis system. Growth of uterine **tissue** was based on
cellular content of DNA and RNA, RNA:DNA, and protein: DNA.
Results-Significant differences in vascular **density** were not
observed in the endometrium of uteri obtained from mares euthanatized
during the follicular or luteal phase of the estrous cycle, regardless of
whether endometrial **classification** of degree of fibrosis was
considered. There was a 3-fold increase in amount of DNA and RNA of
endometrial **cells** in the follicular phase when compared to
myometrium. Hypertrophy of endometrial **tissue** during the luteal
phase was reflected by a significant increase in **cell** protein
content and protein:DNA. Conclusions and Clinical Relevance-Endometrial
growth of vascular **tissues** during the estrous cycle may be
coordinated with development of nonvascular **tissue**. Estrogen and
progesterone may play a role in regulation of uterine growth and
angiogenesis.

L5 ANSWER 2 OF 16 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 2
AN 2000:423342 BIOSIS
DN PREV200000423342
TI Vascular architecture and hypoxic profiles in human head and neck squamous
cell carcinomas.
AU Wyffels, Kiem (1); Kaanders, Jham; Rijken, P. F. J. W.; Bussink, J.; van
den Hoogen, F. J. A.; Marres, Ham; de Wilde, P. C. M.; Raleigh, J. A.; van
der Kogel, A. J.
CS (1) Institute of Radiotherapy, University of Nijmegen, 6500 HB, Nijmegen

Netherlands
 SO British Journal of Cancer, (September, 2000) Vol. 83, No. 5, pp. 674-683.
 print.
 ISSN: 0007-0920.

DT Article
 LA English
 SL English
 AB Tumour oxygenation and vasculature are determinants for radiation treatment outcome and prognosis in patients with squamous cell carcinomas of the head and neck. In this study we visualized and quantified these factors which may provide a predictive tool for new treatments. Twenty-one patients with stage III-IV squamous cell carcinomas of the head and neck were intravenously injected with pimonidazole, a bioreductive hypoxic marker. Tumour biopsies were taken 2 h later. Frozen tissue sections were stained for vessels and hypoxia by fluorescent immunohistochemistry. Twenty-two sections of biopsies of different head and neck sites were scanned and analysed with a computerized image analysis system. The hypoxic fractions varied from 0.02 to 0.29 and were independent from T- and N-classification, localization and differentiation grade. No significant correlation between hypoxic fraction and vascular density was observed. As a first attempt to categorize tumours based on their hypoxia profile, three different hypoxia patterns are described. The first category comprised tumours with large hypoxic, but viable, areas at distances even greater than 200 µm from the vessels. The second category showed a typical band-like distribution of hypoxia at an intermediate distance (50-200 µm) from the vessels with necrosis at greater distances. The third category demonstrated hypoxia already within 50 µm from the vessels, suggestive for acute hypoxia. This method of multiparameter analysis proved to be clinically feasible. The information on architectural patterns and the differences that exist between tumours can improve our understanding of the tumour micro-environment and may in the future be of assistance with the selection of (oxygenation modifying) treatment strategies.

L5 ANSWER 3 OF 16 MEDLINE
 AN 2000511183 MEDLINE
 DN 20518416 PubMed ID: 11064813
 TI Validation of nuclear texture, density, morphometry and tissue syntactic structure analysis as prognosticators of cervical carcinoma.
 AU Weyn B; Tjalma W; Van De Wouwer G; Van Daele A; Scheunders P; Jacob W; Van Marck E
 CS Department of Obstetrics and Gynecology, University of Antwerp, Wilrijk, Belgium.
 SO ANALYTICAL AND QUANTITATIVE CYTOLOGY AND HISTOLOGY, (2000 Oct) 22 (5) 373-82.
 Journal code: ACQ. ISSN: 0884-6812.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 (VALIDATION STUDIES)
 LA English
 FS Priority Journals
 EM 200102
 ED Entered STN: 20010322
 Last Updated on STN: 20010322
 Entered PubMed: 20010131
 Entered Medline: 20010215

AB OBJECTIVE: To evaluate the performance of karyometry and histometry in the prediction of survival, recurrence and response of early-stage invasive cervical carcinoma. STUDY DESIGN: Nuclear morphometry, chromatin texture and tissue architecture (characterized by syntactic structure analysis) were measured using a semiautomated image analysis system on 46 cases of Feulgen-stained tissue sections. The performance of the features was compared to that of clinical features, reported to be the best prognosticators until now, such as age, lympho-vascular permeation, histologic type, stage and grade. A K nearest neighbor classifier was used for classification. RESULTS: In the prediction of three-year survival, recurrence and response, syntactic

structure analysis proved to be the best performer. **Classification** rates were, respectively 100%, 94.4% and 94.5%. In all **classifications**, karyometric and histometric features outperformed clinical features. In general, the best performing features described differences in second-order population statistics (standard deviations). **CONCLUSION:** The results show that a quantitative analysis based on nuclear morphology, chromatin texture and histology can be considered an excellent aid in the prognosis of invasive cervical carcinoma. The measurements are not hampered by the need to undertake complete resections and are suited to daily practice when implemented in a semiautomated **image analysis** system.

L5 ANSWER 4 OF 16 SCISEARCH COPYRIGHT 2001 ISI (R)
 AN 2000:819993 SCISEARCH
 GA The Genuine Article (R) Number: 367JE
 TI Specific changes of chromatin structure in nuclei of normal epithelium adjacent to laryngeal squamous **cell** carcinoma - A preliminary study of 82 cases
 AU Dreyer T (Reprint); Knoblauch I; Garner D; Doudkine A; MacAulay C; Palcic B; Popella C
 CS UNIV GIESSEN, INST PATHOL, LANGHANSSTR 10, D-35392 GIESSEN, GERMANY (Reprint); BRITISH COLUMBIA CANC AGCY, CANC IMAGING DEPT, VANCOUVER, BC V5Z 4E6, CANADA; UNIV GIESSEN, DEPT OTORHINOLARYNGOL, D-35392 GIESSEN, GERMANY
 CYA GERMANY; CANADA
 SO ANALYTICAL CELLULAR PATHOLOGY, (OCT 2000) Vol. 20, No. 2-3, pp. 141-150. Publisher: IOS PRESS, NIEUWE HEMWEG 6B, 1013 BG AMSTERDAM, NETHERLANDS. ISSN: 0921-8912.
 DT Article; Journal
 FS LIFE; CLIN
 LA English
 REC Reference Count: 47
 ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS
 AB The aim of this study was to confirm the existence of specific nuclear texture feature alterations of histologically normal epithelial borders nearby invasive laryngeal cancer (NC).
 Paraffin sections of NC and of chronic inflammations unrelated to cancer (CI) were analysed for nuclear texture and for integrated optical **density** (IOD-index) and were compared to normal epithelium of patients without evidence of cancer (NE). Several discriminant functions based on nuclear texture features were trained to separate different subgroups. As the most important result, specific nuclear texture feature shifts were only found in NC with high-**density** lymphocytic stroma infiltrate (NC+). **Classification** of nuclei of NE versus NC+ was correct in 70%. The same classifier was correct in only 58% when nuclei of NE were classified versus CI. We also found lower values of IOD-Index within the NC+ group when compared to NE (p <0.001).

L5 ANSWER 5 OF 16 MEDLINE
 AN 2000024271 MEDLINE
 DN 20024271 PubMed ID: 10560481
 TI Distinguishing cortical adrenal gland adenomas from carcinomas by their quantitative nuclear features.
 AU Scarpelli M; Montironi R; Mazzucchelli R; Thompson D; Bartels P H
 CS Department of Pathology, University of Ancona, Italy.
 NC R 35 CA 53877 (NCI)
 SO ANALYTICAL AND QUANTITATIVE CYTOLOGY AND HISTOLOGY, (1999 Apr) 21 (2) 131-8.
 Journal code: ACQ; 8506819. ISSN: 0884-6812.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199911
 ED Entered STN: 20000111
 Last Updated on STN: 20000111
 Entered Medline: 19991124
 AB **OBJECTIVE:** To explore data from a set of cases of adrenal cortical adenomas with different endocrine syndromes and carcinomas to determine

whether quantitative **image analysis** of nuclear features might be used to separate the groups. **STUDY DESIGN:** Fifteen adrenal cortical tumors in which clinical information and optimally preserved, paraffin-embedded **tissue** were available were used. There were 10 adenomas and 5 carcinomas. Among the adenomas, five were associated with primary hyperaldosteronism (Conn's syndrome) and five with Cushing's syndrome. Five-micrometer-thick sections were stained with hematoxylin and eosin. In each case 50 nuclei were measured, and a number of morphometric and densitometric features were extracted. The data were subjected to multivariate analysis. **RESULTS:** Quantitative analysis showed that nuclei from adrenal carcinomas are larger than those from adenomas. Total optical **density** (OD) had a near-diploid distribution in the adenomas, while it was clearly aneuploid in the carcinomas. The pixel OD histograms were almost identical for all groups. Differences in nuclear chromatin texture were found between adenomas and carcinomas and also between the two adenoma categories. Multivariate analysis showed good discrimination between carcinomas and adenomas (Wilks lambda = .628, $P < .0001$) and between adenomas. However, based on Bayesian decision boundaries, 20-25% of carcinoma nuclei could be expected to be in the range of adenoma, and about 12% of Cushing's adenoma nuclei and 15% of Conn's adenoma nuclei would be classified as carcinoma. **CONCLUSION:** Computer-assisted analysis of nuclear characteristics proved useful in identifying and describing differences between groups of tumors arising in the adrenal cortex.

L5 ANSWER 6 OF 16 MEDLINE
AN 97133476 MEDLINE
DN 97133476 PubMed ID: 8978872
TI Breast carcinoma. Correlations between visual diagnostic criteria for histologic grading and features of **image analysis**.
AU Tuzek H V; Fritz P; Schwarzmann P; Wu X; Mahner G
CS Department of Pathology, Marienhospital and Robert Bosch Hospital, Stuttgart, Germany.
SO ANALYTICAL AND QUANTITATIVE CYTOLOGY AND HISTOLOGY, (1996 Dec) 18 (6) 481-93.
Journal code: ACQ; 8506819. ISSN: 0884-6812.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 199703
ED Entered STN: 19970327
Last Updated on STN: 19970327
Entered Medline: 19970320
AB **OBJECTIVE:** To investigate the relevance of **image analysis** for grading breast carcinomas. **STUDY DESIGN:** The results of histologic grading were correlated with 18 features of **image analysis**, including SD. "Simple" characteristics, like area and perimeter, shape indices, optical **density** and textural features of nuclei from cancer **cells**, were analyzed. Hematoxylin-eosin-stained **tissue** sections of 67 cancer specimens were routinely used for the study. **RESULTS:** We found statistically significant correlations between overall histologic grading and the sum of its subscores and features of **image analysis**, especially nuclear area, nuclear perimeter and the diameter of the circumscribing circle (diametercirc), including their SDs. The visually and therefore subjectively assessed subscore of the nuclear pleomorphism of histologic grading significantly correlated with the features of **image analysis**, like nuclear area, nuclear perimeter, diametercirc, integrated optical **density** and correlation (and their SDs). There were significant relationships between the absolute numbers of mitoses per 10 high-power fields and nuclear area, nuclear perimeter and diametercirc (and their SDs). We did not observe a significant correlation between the subscore of tubule formation of histologic grading and any of the features of the **image analysis** studied. Furthermore, the correlations between the features of **image analysis** and the subscores of the visual histologic grading system were analyzed with respect to each other. The subscore of nuclear pleomorphism of histologic grading correlated best with overall grading (r

= .72), whereas no significant correlation could be found between the subscores of nuclear pleomorphism and mitotic activity. CONCLUSION: Image analysis provides objectivity and reproducibility to the grading of breast carcinomas and thus could contribute to more individualized prognostication of the disease.

L5 ANSWER 7 OF 16 BIOSIS COPYRIGHT 2001 BIOSIS
AN 1996:327815 BIOSIS
DN PREV199699050171
TI Sample preparation and in situ hybridization techniques for automated molecular cytogenetic analysis of white blood cells.
AU Van De Rijke, Frans M.; Vrolijk, Hans; Sloos, Willem; Tanke, Hans J.; Raap, Anton K. (1)
CS (1) Dep. Cytochem. Cytometry, Leiden Univ., Wassenaarseweg 72, NL 2333 AL Leiden Netherlands
SO Cytometry, (1996) Vol. 24, No. 2, pp. 151-157.
ISSN: 0196-4763.
DT Article
LA English
AB With the advent of in situ hybridization techniques for the analysis of chromosome copy number or structure in interphase cells, the diagnostic and prognostic potential of cytogenetics has been augmented considerably. In theory, the strategies for detection of cytogenetically aberrant cells by in situ hybridization are simple and straightforward. In practice, however, they are fallible, because false classification of hybridization spot number or patterns occurs. When a decision has to be made on molecular cytogenetic normalcy or abnormality of a cell sample, the problem of false classification becomes particularly prominent if the fraction of aberrant cells is relatively small. In such mosaic situations, often gt 200 cells have to be evaluated to reach a statistical sound figure. The manual enumeration of in situ hybridization spots in many cells in many patient samples is tedious. Assistance in the evaluation process by automation of microscope functions and image analysis techniques is, therefore, strongly indicated. Next to research and development of microscope hardware, camera technology, and image analysis, the optimization of the specimen for the (semi)automated microscopic analysis is essential, since factors such as cell density, thickness, and overlap have dramatic influences on the speed and complexity of the analysis process. Here we describe experiments that have led to a protocol for blood cell specimen that results in microscope preparations that are well suited for automated molecular cytogenetic analysis.

L5 ANSWER 8 OF 16 SCISEARCH COPYRIGHT 2001 ISI (R)
AN 96:354409 SCISEARCH
GA The Genuine Article (R) Number: UH687
TI METHODOLOGICAL ASPECTS OF USING DECISION TREES TO CHARACTERIZE LEIOMYOMATOUS TUMORS
AU DECAESTECKER C; REMMELINK M; SALMON I; CAMBY I; GOLDSCHMIDT D; PETEIN M; VANHAM P; PASTEELS J L; KISS R (Reprint)
CS FREE UNIV BRUSSELS, FAC MED, HISTOL LAB, 808 ROUTE LENNIK, B-1070 BRUSSELS, BELGIUM (Reprint); FREE UNIV BRUSSELS, FAC MED, HISTOL LAB, B-1070 BRUSSELS, BELGIUM; FREE UNIV BRUSSELS, ERASME HOSP, INST INTERDISCIPLINARY RES & DEV ARTIFICIAL INTEL, B-1050 BRUSSELS, BELGIUM; FREE UNIV BRUSSELS, ERASME HOSP, FAC MED, HISTOL LAB, B-1050 BRUSSELS, BELGIUM; FREE UNIV BRUSSELS, ERASME HOSP, DEPT PATHOL, B-1050 BRUSSELS, BELGIUM; FREE UNIV BRUSSELS, ERASME HOSP, DEPT PLAST SURG, B-1050 BRUSSELS, BELGIUM; FREE UNIV BRUSSELS, FAC SCI APPL, DEPT DIGITAL & LOG SYST, BRUSSELS, BELGIUM; J BORDET INST, DEPT PATHOL, BRUSSELS, BELGIUM
CYA BELGIUM
SO CYTOMETRY, (01 MAY 1996) Vol. 24, No. 1, pp. 83-92.
ISSN: 0196-4763.
DT Article; Journal
FS LIFE
LA ENGLISH
REC Reference Count: 44
ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS
AB The aim of the present work is to present the potential uses of a

classification technique labeled the "decision tree" for tumor characterisation when used with a large number of features. The decision tree technique enables multifeature logical **classification** rules to be produced by determining discriminatory values for each feature selected. In this report, we propose a methodology that used decision trees to compare and evaluate the information contributed by different types of features for tumor characterisation. This methodology is able to produce a set of hypotheses related to a diagnosis and or prognosis problem. For example, hypotheses can be produced (on the basis of a set of descriptive features) to explain why tumor cases belong to a given histopathological group. To illustrate our purpose, this methodology was applied to the difficult problem of leiomyomatous tumour diagnosis. The aim was to illustrate what kind of diagnostic information can be extracted from a sample data set including 23 smooth muscle tumors (14 benign leiomyomas and 9 malignant leiomyosarcomas) described by a large set of computer-assisted, microscope-generated features. Three groups of features were used relating to: (1) ploidy level determination (10 features), (2) quantitative chromatin pattern description (15 features), and (3) immunohistochemically related antigen specificities (6 features). All these features were quantified by digital **cell image analysis**. The results suggest that an objective distinction between leiomyomas and leiomyosarcomas can be established by means of simple logical rules depending on only a few features among which the immunohistochemically revealed antigen expression of desmin plays a preponderant part. One of the combinations of features proposed by the methodology is interesting for pathologists, because it includes two features describing the appearance of a nucleus in terms of chromatin distribution homogeneity and **density**, two features widely used by pathologists in tumor-grading systems. (C) 1996 Wiley-Liss, Inc.

L5 ANSWER 9 OF 16 SCISEARCH COPYRIGHT 2001 ISI (R)

AN 96:869188 SCISEARCH

GA The Genuine Article (R) Number: VU004

TI Interlaboratory comparison of DNA **image analysis**

AU Thunnissen F B J M (Reprint); Ellis I O; Jutting U

CS UNIV LIMBURG, UNIV HOSP MAASTRICHT, DEPT PATHOL, POB 5800, NL-6202 AZ MAASTRICHT, NETHERLANDS (Reprint); CITY HOSP, DEPT HISTOPATHOL, NOTTINGHAM, ENGLAND; GSF MUNICH, INST PATHOL, MUNICH, GERMANY

CYA NETHERLANDS; ENGLAND; GERMANY

SO ANALYTICAL CELLULAR PATHOLOGY, (OCT 1996) Vol. 12, No. 1, pp. 13-24.

Publisher: ELSEVIER SCI IRELAND LTD, CUSTOMER RELATIONS MANAGER, BAY 15, SHANNON INDUSTRIAL ESTATE CO, CLARE, IRELAND.

ISSN: 0921-8912.

DT Article; Journal

FS LIFE; CLIN

LA English

REC Reference Count: 15

ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS

AB Interlaboratory quality assurance studies have been conducted for DNA flow cytometry, but not for DNA **image analysis** systems. The purpose of this study was to investigate if concordance of DNA **image analysis** systems existed with respect to **classification** and staining of standardized material. In three separate rounds, human liver **cells** were measured randomly by means of the image cytometry system present in each participating laboratory. The features integrated optical **density** (IOD) and AREA were reported. The relationship between the coefficient of variation (CV) of the 2c and 4c peak were compared with three models. In the three rounds the number of participating laboratories was 11, 14 and 11, respectively. Sequential plotting of normalized IOD values yielded useful information about intra-measurement variation. Comparison of measurements in specimens stained in the participating and central laboratory revealed similar CV values. In general, the precision of the instruments, expressed as the 4c/2c and 8c/2c ratios was good. The accuracy of the different laboratories expressed as the CV of IOD for the three rounds varied from 2-17%. The relation of the CVs of the 2c and the 4c peaks was best fit with the model of the addition of two normal distributions. We conclude that interlaboratory comparison of DNA measurements performed on different instruments is certainly feasible and could facilitate improvement in

quality standards.

L5 ANSWER 10 OF 16 MEDLINE
AN 96065086 MEDLINE
DN 96065086 PubMed ID: 7485398
TI Phenotypic analysis of pulmonary perivascular mononuclear infiltrates that occur as a direct result of acute lethal graft-versus-host disease describes the onset of interstitial pneumonitis.
CM Erratum in: Am J Pathol 1996 Feb;148(2):678
AU Workman D L; Clancy J Jr
CS Department of Cell Biology, Neurobiology, and Anatomy, Loyola University Chicago, Maywood, Illinois 60153, USA.
NC AI-12738 (NIAID)
SO AMERICAN JOURNAL OF PATHOLOGY, (1995 Nov) 147 (5) 1350-60.
Journal code: 3RS; 0370502. ISSN: 0002-9440.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Abridged Index Medicus Journals; Priority Journals
EM 199512
ED Entered STN: 19960124
Last Updated on STN: 19980206
Entered Medline: 19951219
AB We recently determined that the sequential development of interstitial pneumonitis and lymphocytic bronchiolitis/bronchitis occurs as a direct result of acute lethal graft-versus-host disease. Interstitial pneumonitis develops before lymphocytic bronchiolitis/bronchitis primarily from the dissemination of perivascular mononuclear infiltrates. We have used the adult, nonirradiated (DA x LEW) F1 hybrid rat in the absence of chemotherapy, immunosuppression, or overt infection to determine the phenotype of infiltrating perivascular mononuclear **cells** throughout acute lethal graft-versus-host disease. F1 animals were intravenously injected with 1×10^6 DA parental lymphoid **cells** /g body weight, which produced 100% morbidity and mortality by day 21. Graft-versus-host disease animals were killed on days 3, 7, 10, 14, and 15 to 21 after injection. Whole left lung lobes were frozen, serially sectioned (4 microns), and incubated with a panel of mouse anti-rat monoclonal antibodies. Labeled antibody **density** was determined by computerized **image analysis**. Perivascular infiltration was observed first for ED1+, OX8+, and W3/25+ **cells**, and then OX41+, W3/13+ and OX19/25+ populations. OX6 was expressed in control **tissues** and at all time points tested. OX12+, OX39+ and MOM/3F12/F2+ **cells** were not quantifiable. The present study has determined that the process of perivascular infiltration was produced through a biphasic influx of OX6+, T-**cell**, and macrophage populations.

L5 ANSWER 11 OF 16 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 3
AN 1995:202135 BIOSIS
DN PREV199598216435
TI Frozen Section Microautoradiography in the Study of Radionuclide Targeting: Application to Indium-111-Oxine-Labeled Leukocytes.
AU Puncher, Matthew R. B.; Blower, Philip J. (1)
CS (1) Nuclear Med. Dep., Kent Canterbury Hosp., Canterbury, CT1 3NH UK
SO Journal of Nuclear Medicine, (1995) Vol. 36, No. 3, pp. 499-505.
ISSN: 0161-5505.
DT Article
LA English
AB The microscopic biodistribution of radioactivity in **tissues** is important in determining microdosimetry. This study addresses the use of frozen section microautoradiography in studying the subcellular distribution of ^{111}In in leukocytes labeled with ^{111}In -oxine. Methods: In conjunction with frozen section microautoradiography, computer **image analysis** methods were applied to the analysis and quantification of leukocyte sections and superimposed autoradiographs. Rapid **cell** fractionation was used to confirm the results. Results: The emulsion (Ilford K2) response was linear over the concentration range investigated (0-33 MBq ml⁻¹). Resolution of radionuclide distribution was better than 2 μm . The autoradiographs

showed no dependence of radiolabel uptake on cell type. **Classification** of all cells into intervals according to grain density suggests an exponential rather than normal distribution, with approximately 50% of cells having little or no radiolabel. In any one sample, cells which were heavily labeled were approximately 10 times more likely to be found in aggregates (60% found in aggregates, mostly neutrophils) than cells which were not heavily labeled (6% found in aggregates); and the grain densities were at least twofold higher over nuclei than over cytoplasm. The last observation was confirmed by the rapid cell fractionation method which showed that approximately 57% of the total radioactivity was bound to nuclei. Conclusion: Frozen section microautoradiography is a practical and reliable approach to determining sub-cellular distribution of ¹¹¹In. The radiolabeling process causes aggregation of neutrophils. Uptake is not significantly dependent on cell type, but only a fraction of cells are appreciably labeled. The radioactive concentration in cell nuclei is at least two-fold higher than in cytoplasm. Microautoradiography can be used to provide distribution data as input into computer models for sub-cellular dosimetry.

L5 ANSWER 12 OF 16 MEDLINE
 AN 95168377 MEDLINE
 DN 95168377 PubMed ID: 7864163
 TI Branching points of renal resistance arteries are enriched in L-type calcium channels and initiate vasoconstriction.
 AU Goligorsky M S; Colflesh D; Gordienko D; Moore L C
 CS Department of Medicine, State University of New York at Stony Brook 11794-8152.
 NC DK-26341 (NIDDK)
 DK-41573 (NIDDK)
 RR-05736 (NCRR)
 SO AMERICAN JOURNAL OF PHYSIOLOGY, (1995 Feb) 268 (2 Pt 2) F251-7.
 Journal code: 3U8; 0370511. ISSN: 0002-9513.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 199503
 ED Entered STN: 19950404
 Last Updated on STN: 19950404
 Entered Medline: 19950320
 AB The morphologic structures responsible for the drop in blood pressure along the preglomerular vasculature are not completely defined. Theoretical and videomicroscopic analyses of nonrenal vascular beds implicate bifurcations of resistance arteries as important sites of hemodynamic regulation. These structures contain pacemaker cells sensitive to calcium channel blockers and appear to initiate vasomotion. In the present study, we examined the possibility of functional diversity of smooth muscle cells along resistance arteries with regard to the density of voltage-gated L-type calcium channels. Staining of microdissected renal resistance arteries with Bodipy-labeled dihydropyridine and analysis by confocal microscopy showed enhanced binding at branching points compared with the distal sites in daughter vessels. Antibodies directed against the alpha 1-subunit of the dihydropyridine-sensitive calcium channels confirmed the enhanced expression of L-type channels predominantly at the sites of bifurcations of renal resistance arteries. Fluorescence digital-image analysis of freshly microdissected branches of cortical radial (interlobular) and arcuate arteries intravitaly labeled with a calcium indicator, fluo 3, identified branching points as initiator sites of depolarization-induced intracellular Ca²⁺ concentration ([Ca²⁺]_i) transients, which propagated along the vascular wall at the rate of 2.0 +/- 0.7 micron/s. Videomicroscopy of blood-perfused rat juxtamedullary resistance arteries showed that branching points exhibit more pronounced contractile responses to KCl-induced depolarization than distal sites along the daughter vessels. Collectively, these results demonstrate that branching points are enriched in L-type calcium channels, a finding that suggests these structures may serve as important regulators of renal

L5 ANSWER 13 OF 16 MEDLINE
AN 95068610 MEDLINE
DN 95068610 PubMed ID: 7526719
TI Computer-assisted **image analysis** of tumor sections for
a new thrombospondin receptor.
AU Arnoletti J P; Albo D; Jhala N; Granick M S; Solomon M P; Atkinson B;
Rothman V L; Tuszynski G P
CS Department of Surgery, Medical College of Pennsylvania, Philadelphia
19129.
SO AMERICAN JOURNAL OF SURGERY, (1994 Nov) 168 (5) 433-6.
Journal code: 3Z4; 0370473. ISSN: 0002-9610.
CY United States
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Abridged Index Medicus Journals; Priority Journals
EM 199412
ED Entered STN: 19950110
Last Updated on STN: 19960129
Entered Medline: 19941207
AB BACKGROUND: A **cell** surface receptor (50 kd) has been recently
identified in malignant **cells** that recognizes the tumor
cell adhesive domain (ie, cysteine-serine-valine-threonine-
cysteine-glycine [CSVTCG]) of thrombospondin (TSP). This CSVTCG-specific
TSP receptor can be considered as a new tumor marker, and its
concentration on the **cell** surface may correlate directly with
the capacity of tumor **cells** to invade and metastasize. MATERIALS
AND METHODS: Six patients with primary, stages III and IV squamous
cell carcinomas of the head and neck were studied. Tumor sections
were specifically stained for this receptor with immunohistochemical
techniques. The stained specimens were then subjected to computer-assisted
image analysis. The area of positive staining and the
heterogeneity of the pattern of staining were compared to peritumoral
angiogenesis and clinical outcome of the patients. RESULTS: The results
indicate that those patients with a high and homogenous positive stain
score (mean +/- standard error [SE] 78 +/- 5%) for the CSVTCG-specific TSP
receptor had high microvessel **density** and died from metastatic
disease within 12 months of initial treatment (correlation coefficients =
0.95 and 1, respectively). Patients with a low and heterogenous positive
stain score for receptor (mean +/- SE 8 +/- 2%; P < 0.001) had low
microvessel counts and remained disease-free for at least 2 years. There
was no relationship between receptor **density** and histologic
classification of the primary tumors. CONCLUSION: The
CSVTCG-specific TSP receptor, quantified through **image**
analysis of immunohistochemical stained **tissue** sections,
is highly predictive of clinical outcome in patients with squamous
cell carcinomas of the head and neck.

L5 ANSWER 14 OF 16 BIOSIS COPYRIGHT 2001 BIOSIS
AN 1993:230606 BIOSIS
DN PREV199395121781
TI Method for counting mitoses by image processing in Feulgen stained breast
cancer sections.
AU Ten Kate, T. K.; Belien, J. A. M.; Smeulders, A. W. M.; Baak, J. P. A. (1)
CS (1) Dep. Pathology, Free Univ. Hosp., de Boelelaan 1117, 1007 MB
Amsterdam, The Netherlands
SO Cytometry, (1993) Vol. 14, No. 3, pp. 241-250.
ISSN: 0196-4763.
DT Article
LA English
AB This study describes an image processing method for the assessment of the
mitotic count in Feulgen-stained breast cancer sections. The segmentation
procedure was optimized to eliminate 95-98% of the nonmitoses, whereas 11%
of the mitoses did not survive the segmentation procedure. Contour
features and optical **density** measurements of the remaining
objects were computed to allow for **classification**. Twelve
specimens were analyzed, nine used to serve as a training set, and three
put aside for later use as independent test set. The fully automatic image

- processing method correctly classified 81% of the mitoses at the specimen level while inserting false positives. The automatic procedure strongly correlated with the interactive counting procedure ($r=0.98$). Although the fully automatic method provided satisfactory results, it is not yet suited for clinical practice. The automated method with an interactive evaluation step gave an accurate reflection of the mitotic count showing an almost perfect correlation with the results of the interactive morphometry ($r=0.998$). Therefore this semiautomated method may be useful as prescreening device.

L5 ANSWER 15 OF 16 MEDLINE

AN 92316470 MEDLINE

DN 92316470 PubMed ID: 1618471

TI Changes of IgG-bearing **cell** populations in the portal tracts of patients with chronic liver disease of viral etiology: an evaluation by immunoperoxidase method and computerized **image analysis**

AU Torgano G; Vecchi M; Arosio E; Santambrogio D; Ronchi G; Annoni G; Tomasini M; Contessini E; de Franchis R

CS Department of Internal Medicine, University of Milano, Italy.

SO HEPATOLOGY, (1992 Jul) 16 (1) 19-23.

Journal code: GBZ; 8302946. ISSN: 0270-9139.

CY United States

DT Journal; Article; (JOURNAL ARTICLE)

LA English

FS Priority Journals

EM 199208

ED Entered STN: 19920815

Last Updated on STN: 19920815

Entered Medline: 19920803

AB Little is known about the distribution of IgG-bearing **cell** subpopulations in normal liver and their possible changes in disease conditions. We developed an immunohistochemical method that proved suitable and accurate for the identification and characterization of IgG-bearing **cells** and their subpopulations in liver specimens. The method uses specific monoclonal antibodies on serial mirror liver sections. We applied this method to four normal liver **tissue** specimens and 25 liver biopsy samples of chronic hepatitis of viral etiology. Only rare IgG-bearing **cells** could be observed in the portal tracts of normal liver specimens. In contrast, a dense infiltrate of such **cells** was seen in liver specimens from patients with chronic viral hepatitis. The **density** of IgG-bearing **cells** in such patients ranged from 6 to 20 **cells** x $10(-4)$ micron² in the different specimens (mean = $11 \times 10(-4)$ micron²). The increase in IgG-bearing **cells** did not appear to be related to the histological diagnosis, to the degree of histological inflammatory activity or to the type of viral infection. The major population of IgG-bearing **cells** consisted of IgG1-positive **cells** (68%); IgG2- (17%), IgG3- (8%) and IgG4 (7%)-bearing **cells** represented only minor fractions. The increased prevalence of IgG1-bearing **cells** observed in chronic hepatitis but not in normal liver specimens suggests that these findings may reflect an activation of antibody production directed toward viral antigens or antigenic structures of self. The identification of the antigenic specificities of the antibodies produced by IgG-bearing **cells** might provide important clues in understanding the pathogenesis of chronic viral hepatitis.

L5 ANSWER 16 OF 16 BIOSIS COPYRIGHT 2001 BIOSIS

AN 1991:392904 BIOSIS

DN BA92:70219

TI ASSESSEMENT OF **CELL** PROLIFERATION ON POROUS MICROCARRIERS BY MEANS OF **IMAGE ANALYSIS**.

AU FORAN D J; CAHN F; EIKENBERRY E F

CS DEP. PATHOL., ROBERT WOOD JOHNSON MED. SCH., UNIV. MED. DENTISTRY NEW JERSEY, PISCATAWAY, N.J. 08854.

SO ANAL QUANT CYTOL HISTOL, (1991) 13 (3), 215-222.

CODEN: AQCHED. ISSN: 0884-6812.

FS BA; OLD

LA English

AB Spherical porous microcarriers (PMCs) made from collagen-glycosaminoglycan crosslinked copolymers have exhibited considerable promise as growth surfaces for the proliferation of anchorage-dependent mammalian cell lines and have demonstrated the ability to entrap anchorage-independent cells. However, quantification of cell growth on PMCs has proved difficult. A method of measuring the proliferation of PMCs, based on image analysis, is presented. Using CV1 and CHO cell lines, samples of PMCs were removed from culture at various times, fixed, embedded and sectioned. The 2 .mu.m sections were stained, photographed and digitized in three colors. A computer program was developed to evaluate digitized PMC cross-sections and to classify pixels as conforming to either background, cytoplasmic, matrix or nuclear parameters, based on a set of classification rules determined by statistical analysis. Growth curves were generated by relating the number of pixels occupied by cellular material to the total number of pixels in the PMC cross-section. The PMCs were found to foster cell proliferation, with cell densities approaching 100% occupancy.

=> d his

(FILE 'HOME' ENTERED AT 20:47:44 ON 22 MAY 2001)

FILE 'BIOSIS, CAPLUS, MEDLINE, SCISEARCH' ENTERED AT 20:48:11 ON 22 MAY 2001

L1	0 S CELL DENISTY AND IMAGE ANALYSIS
L2	0 S CELL DENISTY AND CLASSIFICATION
L3	196 S CELL AND CLASSIFICATION AND IMAGE ANALYSIS AND TISSUE
L4	22 S L3 AND DENSITY
L5	16 DUP REM L4 (6 DUPLICATES REMOVED)

=> log y

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	58.72	58.87

STN INTERNATIONAL LOGOFF AT 21:01:46 ON 22 MAY 2001